



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

### SOLUTIONS OF PROBLEMS.

43. Proposed by HENRY HEATON, M. Sc., Atlantic, Iowa.

In a circle whose radius is  $a$ , chords are drawn through a point distant  $b$  from the center. What is the average length of such chords, (1), if a chord is drawn from every point of the circumference, and (2), if they are drawn through the point at equal angular intervals?

II. Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas.

Let  $AC=CF=a$ ,  $CD=b$ ,  $\angle FDA=\theta$ ,  $\angle FCA=\varphi$ ,  $\tan\theta=m$ .

Then the equation to  $DF$  is,  $y=mx+mb$ .....(1).

The equation to the circle is,  $x^2+y^2=a^2$ .....(2).

From (1) and (2) we easily get  $EF=2\sqrt{a^2-\frac{m^2b^2}{1+m^2}}$ ,

$\therefore EF=2\sqrt{a^2-b^2\sin^2\theta}$ . But  $\sin\theta=\frac{a\sin\varphi}{\sqrt{a^2+b^2+2ab\cos\varphi}}$ .

$\therefore EF=\frac{2(a^2+ab\cos\varphi)}{\sqrt{a^2+b^2+2ab\cos\varphi}}=\frac{2(a^2+ab-2ab\sin^2\frac{1}{2}\varphi)}{\sqrt{(a+b)^2-4ab\sin^2\frac{1}{2}\varphi}}$ .

The limits of  $\varphi$  for  $b>a$ , are 0 and  $\frac{1}{2}\pi+\sin^{-1}(a/b)=2\beta$ .

The limits of  $\varphi$  for  $b<a$ , are 0 and  $\frac{1}{2}\pi+\sin^{-1}(b/a)=2\delta$ .

The limits of  $\theta$  for  $b>a$ , are 0 and  $\sin^{-1}(a/b)=\theta'$ .

The limits of  $\theta$  for  $b<a$ , are 0 and  $\frac{1}{2}\pi$ .


Let  $\Delta$  and  $\Delta_1$  be the average lengths required.

$\therefore \Delta=\int E F d s / \int d s=\int E F d \varphi / \int d \varphi$ ,

$\Delta_1=\int E F d \theta / \int d \theta$ .

I. Let  $\frac{4ab}{(a+b)^2}=e^2$ , and  $\frac{1}{2}\varphi=\gamma$ .

$\therefore d\varphi=2d\gamma$ .



$$\text{Then } EF = \frac{2(a^2 + ab - 2ab\sin^2\gamma)}{(a+b)\sqrt{1-e^2\sin^2\gamma}}.$$

$$\therefore EF = \frac{2a}{\sqrt{1-e^2\sin^2\gamma}} + \frac{2}{e\sqrt{ab}}\sqrt{1-e^2\sin^2\gamma} - \frac{2}{e\sqrt{ab}\sqrt{1-e^2\sin^2\gamma}}.$$

$$\therefore \int EF d\gamma = \frac{2}{e\sqrt{ab}} \{ (ae\sqrt{ab} - 1)F(e, \gamma) + E(e, \gamma) \}.$$

$$\therefore A = \frac{2}{\beta e\sqrt{ab}} \{ (ae\sqrt{ab} - 1)F_0^\beta(e, \gamma) + E_0^\beta(e, \gamma) \}, \quad b > a.$$

$$A = \frac{2}{\delta e\sqrt{ab}} \{ (ae\sqrt{ab} - 1)F_0^\delta(e, \gamma) + E_0^\delta(e, \gamma) \}, \quad b < a.$$

$$\text{II. } A_1 = 2 \int_0^{\theta'} \sqrt{a^2 - b^2 \sin^2 \theta} d\theta / \int_0^{\theta'} d\theta = \frac{2}{\theta'} \int_0^{\theta'} \sqrt{a^2 - b^2 \sin^2 \theta} d\theta.$$

$$\text{Let } \theta = \frac{1}{2}\pi + \lambda. \quad \therefore \theta' - \frac{1}{2}\pi = \lambda \text{ to } -\frac{1}{2}\pi = \lambda.$$

$$\begin{aligned} \therefore A_1 &= \frac{2}{\theta'} \int_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \sqrt{a^2 - b^2 \cos^2 \lambda} d\lambda = \frac{2}{\theta'} \int_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \sqrt{b^2 \sin^2 \lambda - (b^2 - a^2)} d\lambda \\ &= \frac{2\sqrt{(b^2 - a^2)}}{\theta'} \int_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \sqrt{\frac{b^2}{b^2 - a^2} \sin^2 \lambda - 1} d\lambda \\ &= \frac{2\sqrt{(b^2 - a^2)}}{\theta'} H_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \left( \frac{b}{\sqrt{b^2 - a^2}}, \lambda \right), \quad b > a. \end{aligned}$$

$$A_1 = 2 \int_0^{\frac{1}{2}\pi} \sqrt{a^2 - b^2 \sin^2 \theta} d\theta / \int_0^{\frac{1}{2}\pi} db = \frac{4a}{\pi} E_{\frac{1}{2}\pi}^{\frac{1}{2}\pi} \left( \frac{b}{a}, \theta \right), \quad b < a.$$

45. Proposed by J. C. WILLIAMS, Boston, Massachusetts.

At the end of the fifth inning the base ball score stands 7 to 9. What is the probability of winning for either team?

Solution by J. SCHEFFER, A. M., Hagerstown, Maryland.

From the stated score we are able to estimate the respective skill of the two teams, and their respective probabilities of winning the game.

The respective probabilities are  $\frac{7}{18}$  and  $\frac{9}{18}$ . We have now to find the probabilities of either team winning at least 3 games out of 4, granting, of course, 9